


MATTHEW J. DUBIEL M.D. SHOULDER AND ELBOW SURGERY

WWW.MDUBIELMD.COM



KNEE SURGERY

DIET

- BEGIN WITH CLEAR LIQUIDS AND LIGHT FOODS (JELLOS, SOUPS, ETC.)
- PROGRESS TO YOUR NORMAL DIET IF YOU ARE NOT NAUSEATED

WOUND CARE

- MAINTAIN YOUR OPERATIVE DRESSING, YOU CAN LOOSEN BANDAGE IF SWELLING OCCURS
- IT IS NORMAL FOR THE INCISIONS TO BLEED AND SWELL FOLLOWING SURGERY – IF BLOOD SOAKS THROUGH THE BANDAGE, DO NOT BECOME ALARMED – REINFORCE WITH ADDITIONAL DRESSING
- KEEP YOUR DRESSINGS ON UNTIL YOUR APPOINTMENT
- YOU CAN SHOWER TWO DAYS AFTER SURGERY. KEEP YOUR DRESSINGS CLEAN AND DRY,

MEDICATIONS

- YOU MAY HAVE HAD NUMBING MEDICATION INJECTED NEAR YOUR WOUND AND THIS SHOULD WEAR OFF BETWEEN 12 AND 24 HOURS.
- ONLY USE YOUR NARCOTIC PAIN MEDICATION IF NEEDED
- TAKE ASPIRIN ONCE A DAY FOR THREE WEEKS TO DECREASE THE RISK OF BLOOD CLOTS

PRIMARY MEDICATION

- TAKE 1 – 2 TABLETS EVERY 4 – 6 HOURS AS NEEDED MAX OF 12 PILLS PER DAY
- PLAN ON USING IT FOR 5 TO 7 DAYS, DEPENDING ON LEVEL OF PAIN
 - DO NOT TAKE ADDITIONAL TYLENOL (ACETAMINOPHEN) WHILE TAKING NORCO/ PERCOCET
COMMON SIDE EFFECTS OF THE PAIN MEDICATION ARE NAUSEA, DROWSINESS, AND CONSTIPATION – TO DECREASE THE SIDE EFFECTS, TAKE MEDICATION WITH FOOD.
 - IF CONSTIPATION OCCURS, CONSIDER TAKING AN OVER-THE-COUNTER LAXATIVE
 - IF YOU ARE HAVING PROBLEMS WITH NAUSEA AND VOMITING, CONTACT THE OFFICE TO POSSIBLY HAVE YOUR MEDICATION CHANGED
 - DO NOT DRIVE A CAR OR OPERATE MACHINERY WHILE TAKING THE NARCOTIC MEDICATION
 - IBUPROFEN 400-600MG (I.E. ADVIL) MAY BE TAKEN IN BETWEEN THE NARCOTIC PAIN MEDICATION TO HELP SMOOTH OUT THE POST-OPERATIVE 'PEAKS AND VALLEYS', REDUCE OVERALL AMOUNT OF PAIN MEDICATION REQUIRED, AND INCREASE THE TIME INTERVALS BETWEEN NARCOTIC PAIN MEDICATION USAGE.

MATTHEW J. DUBIEL M.D.
SHOULDER AND ELBOW SURGERY

WWW.MDUBIELMD.COM

ACTIVITY

- DO NOT PUT WEIGHT ON YOUR ANKLE
- DO NOT ENGAGE IN ACTIVITIES WHICH INCREASE PAIN/SWELLING OVER THE FIRST 7-10 DAYS FOLLOWING SURGERY
- MAY RETURN TO SEDENTARY WORK ONLY OR SCHOOL 1-2 DAYS AFTER SURGERY, IF PAIN IS TOLERABLE

EMERGENCIES

- CONTACT DR. DUBIEL OR HIS TEAM AT 517-394-3200 IF ANY OF THE FOLLOWING ARE PRESENT:
 - PAINFUL SWELLING OR NUMBNESS
 - UNRELENTING PAIN
 - FEVER (OVER 101° – IT IS NORMAL TO HAVE A LOW GRADE FEVER FOR THE FIRST DAY OR TWO FOLLOWING SURGERY) OR CHILLS
 - REDNESS AROUND INCISIONS
 - COLOR CHANGE IN WRIST, HAND OR LOWER EXTREMITY
 - CONTINUOUS DRAINAGE OR BLEEDING FROM INCISION (A SMALL AMOUNT OF DRAINAGE IS EXPECTED)
 - DIFFICULTY BREATHING
 - EXCESSIVE NAUSEA/VOMITING
- IF YOU HAVE AN EMERGENCY AFTER OFFICE HOURS OR ON THE WEEKEND, CALL 517-394-3200 AND YOU WILL BE CONNECTED TO OUR PAGE SERVICE – THEY WILL CONTACT DR. DUBIEL OR ONE OF HIS PARTNERS IF HE IS UNAVAILABLE. DO NOT CALL THE HOSPITAL OR SURGICENTER.
- IF YOU HAVE AN EMERGENCY THAT REQUIRES IMMEDIATE ATTENTION, PROCEED TO THE NEAREST EMERGENCY ROOM.
- IF YOU DO NOT ALREADY HAVE A POSTOPERATIVE APPOINTMENT SCHEDULED, PLEASE CONTACT THE SCHEDULING OFFICE DURING NORMAL OFFICE HOURS (517-394-3200)
- THIS WEB SITE ALSO INCLUDES MORE POSTOPERATIVE INFORMATION FOR SPECIFIC PROCEDURES, WHICH MAY BE HELPFUL FOR YOUR RECOVERY. (COMING SOON)