


# MATTHEW J. DUBIEL M.D. SHOULDER AND ELBOW SURGERY



## CAPSULAR RELEASE

	RANGE OF MOTION	SLING	EXERCISES
PHASE I 0-4 WEEKS	PASSIVE TO ACTIVE RANGE AS TOLERATED	0-2 WEEKS: WORN FOR COMFORT ONLY  2-4 WEEKS: DISCONTINUE	0-2 WEEKS: INITIATE OUTPATIENT PT ACCORDING TO RX  AGGRESSIVE PROM AND CAPSULAR STRETCHING*; CLOSED CHAIN SCAPULA  2-4 WEEKS: CONTINUE CAPSULAR STRETCHING: PROM, JOINT MOBILIZATION TO MAX TOLERANCE**  DELTOID, CUFF ISOMETRICS, BEGIN SCAPULAR PROTRACTION/ RETRACTION
PHASE II 4-8 WEEKS	INCREASE AS TOLERATED TO FULL	NONE	ADVANCE ISOMETRICS, ROTATOR CUFF AND DELTOID*  ADVANCE TO THERABANDS, DUMBBELLS AS TOLERATED**  CONTINUE CAPSULAR STRETCHING AND PROM
PHASE III 8-16 WEEKS	PROGRESS TO FULL MOTION WITHOUT DISCOMFORT	NONE	ADVANCE STRENGTHENING AS TOLERATED BEGIN ECCENTRICALLY RESISTED MOTIONS AND CLOSED CHAIN ACTIVITIES  ADVANCE TO SPORT AND FULLY ACTIVITY AS TOLERATED AFTER 12 WEEKS